

Datasheet for ABIN7636753 **anti-CXCR1 antibody**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	CXCR1
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CXCR1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Interleukin 8 Receptor Alpha (IL8Ra)
Immunogen:	RPA019Mu01Recombinant Interleukin 8 Receptor Alpha (IL8Ra)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against IL8Ra. It has been selected for its ability to recognize IL8Ra in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	CXCR1
Alternative Name:	IL8Ra (CXCR1 Products)

Target Details

Background: CD181, CDw128a, CXCR1, CKR1, CMKAR1, IL8R1, IL8RBA, IL8-RA, Chemokine C-X-C motif Receptor 1, High Affinity Interleukin-8 Receptor A

UniProt: [Q810W6](#)

Pathways: [cAMP Metabolic Process](#)

Application Details

Application Notes: Western blotting: 0.5-2 µg/mL Immunohistochemistry: 5-20 µg/mL Immunocytochemistry: 5-20 µg/mL Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 500 µg/mL

Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.