

Datasheet for ABIN7636823

anti-Cholecystokinin antibody



Overview

Quantity:	100 μL
Target:	Cholecystokinin (CCK)
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Cholecystokinin antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Target:

Alternative Name:

Froduct Details	
Purpose:	Polyclonal Antibody to Cholecystokinin 8 (CCK8)
Immunogen:	CPB044Hu210VA Conjugated Cholecystokinin 8 (CCK8)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CCK8. It has been selected for its ability to recognize CCK8 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

Cholecystokinin (CCK)

CCK8 (CCK Products)

Target Details

Storage Comment:

Target Details	
Background:	Sincalide, Cholecystokinin 8, Octapeptide
Pathways:	TCR Signaling, Activation of Innate immune Response, Cellular Response to Molecule of Bacterial Origin, Positive Regulation of Immune Effector Process, Positive Regulation of Endopeptidase Activity, Toll-Like Receptors Cascades, Feeding Behaviour
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	500 μg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C

detectable loss of activity. Avoid repeated freeze-thaw cycles.

Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without