

#### Datasheet for ABIN7636993

# anti-Claudin 18 antibody



#### Overview

Quantity:	100 μL
Target:	Claudin 18 (CLDN18)
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Claudin 18 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

#### **Product Details**

Target:

Alternative Name:

Product Details	
Purpose:	Polyclonal Antibody to Claudin 18 (CLDN18)
Immunogen:	RPB100Hu01Recombinant Claudin 18 (CLDN18)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against CLDN18. It has been selected for its ability to recognize CLDN18 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

Claudin 18 (CLDN18)

CLDN18 (CLDN18 Products)

### **Target Details**

Background:	SFTP-J, SFTPJ, SP-J, SPJ, Surfactant Protein J, Surfactant Associated Protein J
UniProt:	P56856
Pathways:	Cell-Cell Junction Organization, Hepatitis C

### **Application Details**

Application Notes:	Western blotting: 0.01-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-
	20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	0.41 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.