

### Datasheet for ABIN7637234

# anti-M-CSF/CSF1 antibody



_					
	W	0	rv	10	W

Quantity:	100 μL
Target:	M-CSF/CSF1 (CSF1)
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This M-CSF/CSF1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

### **Product Details**

Purpose:	Polyclonal Antibody to Colony Stimulating Factor 1, Macrophage (MCSF)
Immunogen:	RPA090Mu02Recombinant Colony Stimulating Factor 1, Macrophage (MCSF)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against MCSF. It has been selected for its ability to recognize MCSF in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

Target:	M-CSF/CSF1 (CSF1)
Alternative Name:	MCSF (CSF1 Products)

## **Target Details**

Background:	CSF1, M-CSF, Lanimostim, Processed macrophage colony-stimulating factor 1	
UniProt:	P07141	
Pathways:	RTK Signaling	

## **Application Details**

Application Notes:	Western blotting: 0.5-2 μg/mL,lmmunohistochemistry: 5-20 μg/mL,lmmunocytochemistry: 5-	
	20 μg/mL,Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	

# Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.