

Datasheet for ABIN7637679 **anti-CYR61 antibody**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	CYR61
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CYR61 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Monoclonal Antibody to Cysteine Rich Protein, Angiogenic Inducer 61 (CYR61)
Immunogen:	RPG313Hu01 Recombinant Cysteine Rich Protein, Angiogenic Inducer 61 (CYR61)
Clone:	C4
Specificity:	The antibody is a mouse monoclonal antibody raised against CYR61. It has been selected for its ability to recognize CYR61 in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography

Target Details

Target:	CYR61
Alternative Name:	CYR61 (CYR61 Products)

Target Details

Background:	CCN1, GIG1, IGFBP10, CCN family member 1, Insulin-like growth factor-binding protein 10
UniProt:	O00622
Pathways:	Positive Regulation of Endopeptidase Activity , Growth Factor Binding

Application Details

Application Notes:	Western blotting: 0.2-2 µg/mL, 1:500-5000 Immunohistochemistry: 5-20 µg/mL, 1:50-200 Immunocytochemistry: 5-20 µg/mL, 1:50-200 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin, Sodium azide
Precaution of Use:	This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.