

Datasheet for ABIN7637955

anti-DGAT2 antibody



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	DGAT2
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DGAT2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Polyclonal Antibody to Diacylglycerol-O-Acyltransferase Homolog 2 (DGAT2)
Immunogen:	RPJ272Hu01 Recombinant Diacylglycerol O-Acyltransferase Homolog 2 (DGAT2)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against DGAT2. It has been selected for its ability to recognize DGAT2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	DGAT2
Alternative Name:	DGAT2 (DGAT2 Products)

Target Details

Background: ARAT, Acyl-CoA retinol O-fatty-acyltransferase, Diglyceride acyltransferase 2

UniProt: [Q96PD7](#)

Application Details

Application Notes: Western blotting: 0.2-2 µg/mL, 1:250-2500 Immunohistochemistry: 5-20 µg/mL, 1:25-100
Immunocytochemistry: 5-20 µg/mL, 1:25-100 Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 0.5 mg/mL

Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.

Preservative: ProClin, Sodium azide

Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.