

Datasheet for ABIN7638144

anti-DNAH11 antibody



Overview

Quantity:	100 μL
Target:	DNAH11
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DNAH11 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Target:

Alternative Name:

Purpose:	Polyclonal Antibody to Dynein, Axonemal, Heavy Chain 11 (DNAH11)
Immunogen:	RPJ201Hu02Recombinant Dynein, Axonemal, Heavy Chain 11 (DNAH11)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against DNAH11. It has been selected for its ability to recognize DNAH11 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

DNAH11

DNAH11 (DNAH11 Products)

Target Details

ranger Details		
Background:	DNAHBL, DNAHC11, DNHBL, DPL11, Dynein, Heavy Chain Beta-Like, Axonemal beta dynein heavy chain 11, Ciliary dynein heavy chain 11	
UniProt:	Q96DT5	
Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-20 μg/mL,Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	0.5 mg/mL	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	ProClin, Sodium azide	
Precaution of Use:	This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	