

Datasheet for ABIN7638333

anti-EZH2 antibody



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	EZH2
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EZH2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Enhancer Of Zeste Homolog 2 (EZH2)
Immunogen:	RPK073Hu02Recombinant Enhancer Of Zeste Homolog 2 (EZH2)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against EZH2. It has been selected for its ability to recognize EZH2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	EZH2
Alternative Name:	EZH2 (EZH2 Products)

Target Details

Background:	ENX-1, KMT6, KMT6A, Histone-lysine N-methyltransferase EZH2, Lysine N-methyltransferase 6
UniProt:	Q15910
Pathways:	Retinoic Acid Receptor Signaling Pathway , Regulation of Muscle Cell Differentiation

Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	500 µg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.