

Datasheet for ABIN7638906
anti-FGF7 antibody



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	FGF7
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This FGF7 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Monoclonal Antibody to Fibroblast Growth Factor 7 (FGF7)
Immunogen:	RPA636Hu01 Recombinant Fibroblast Growth Factor 7 (FGF7)
Specificity:	The antibody is a mouse monoclonal antibody raised against FGF7. It has been selected for its ability to recognize FGF7 in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography

Target Details

Target:	FGF7
Alternative Name:	FGF7 (FGF7 Products)
Background:	KGF, HBGF7, Keratinocyte Growth Factor, Heparin-binding growth factor 7

Target Details

UniProt:	P21781
Pathways:	RTK Signaling , Fc-epsilon Receptor Signaling Pathway , EGFR Signaling Pathway , Neurotrophin Signaling Pathway

Application Details

Application Notes:	Western blotting: 0.2-2 µg/mL, 1:500-5000 Immunohistochemistry: 5-20 µg/mL, 1:50-200 Immunocytochemistry: 5-20 µg/mL, 1:50-200 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin, Sodium azide
Precaution of Use:	This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.