

### Datasheet for ABIN7639177

# anti-GPER antibody



#### Overview

Quantity:	100 μL
Target:	GPER
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GPER antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

## **Product Details**

Target:

Alternative Name:

**GPER** 

**GPER (GPER Products)** 

Purpose:	Polyclonal Antibody to G Protein Coupled Estrogen Receptor 1 (GPER)
Immunogen:	RPG045Mu01Recombinant G Protein Coupled Estrogen Receptor 1 (GPER)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against GPER. It has been selected for its ability to recognize GPER in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

## Target Details

Background:	GPR30, CMKRL2, DRY12, FEG-1, GPCR-Br, LERGU, LyGPR, Membrane estrogen receptor, Chemoattractant receptor-like 2, Flow-induced endothelial G-protein coupled receptor 1
UniProt:	Q8BMP4
Pathways:	EGFR Signaling Pathway, Positive Regulation of Peptide Hormone Secretion, Intracellular Steroid Hormone Receptor Signaling Pathway, Steroid Hormone Mediated Signaling Pathway, Carbohydrate Homeostasis, cAMP Metabolic Process, Regulation of G-Protein Coupled Receptor Protein Signaling, Positive Regulation of Endopeptidase Activity, Regulation of Carbohydrate Metabolic Process

	Steroid Hormone Receptor Signaling Pathway, Steroid Hormone Mediated Signaling Pathway,
	Carbohydrate Homeostasis, cAMP Metabolic Process, Regulation of G-Protein Coupled
	Receptor Protein Signaling, Positive Regulation of Endopeptidase Activity, Regulation of
	Carbohydrate Metabolic Process
Application Details	
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100
	Immunocytochemistry: 5-20 µg/mL,1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin, Sodium azide
Precaution of Use:	This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANCES
	which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.