

Datasheet for ABIN7639676

anti-CD236/GYPC antibody



Overview

Quantity:	100 μL
Target:	CD236/GYPC (GYPC)
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CD236/GYPC antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Target:

Alternative Name:

Troduct Details	
Purpose:	Polyclonal Antibody to Glycophorin C (GYPC)
Immunogen:	RPB711Mu01Recombinant Glycophorin C (GYPC)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against GYPC. It has been selected for its ability to recognize GYPC in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

CD236/GYPC (GYPC)

GYPC (GYPC Products)

Target Details

l arget Details	
Target Type:	Viral Protein
Background:	CD236, CD236R, GYP-C, PAS-2, GPC, GYPD, Ge, Glycoprotein Beta, Glycoconnectin, Gerbich
	Blood Group, Glycophorin-D, Sialoglycoprotein D
UniProt:	Q78HU7
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-
	20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without

detectable loss of activity. Avoid repeated freeze-thaw cycles.