

Datasheet for ABIN7639944 **anti-HAUS7 antibody**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	HAUS7
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HAUS7 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)

Product Details

Purpose:	Polyclonal Antibody to HAUS Augmin Like Complex Subunit 7 (HAUS7)
Immunogen:	RPM623Hu01 Recombinant HAUS Augmin Like Complex Subunit 7 (HAUS7)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against HAUS7. It has been selected for its ability to recognize HAUS7 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	HAUS7
Alternative Name:	HAUS7 (HAUS7 Products)

Target Details

Background: UCHL5IP, UIP1, UCHL5 Interacting Protein, UCH37 Interacting Protein 1, 26S Proteasome-Associated UCH Interacting Protein 1, X-linked protein STS1769

UniProt: [Q99871](#)

Application Details

Application Notes: Western blotting: 0.01-3 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunofluorescence: 5-20 µg/mL, Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 0.5 mg/mL

Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.