

Datasheet for ABIN7640266
anti-HMGB3 antibody



[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	HMGB3
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HMGB3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Polyclonal Antibody to High Mobility Group Box Protein 3 (HMGB3)
Immunogen:	RPD773Hu01Recombinant High Mobility Group Box Protein 3 (HMGB3)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against HMGB3. It has been selected for its ability to recognize HMGB3 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	HMGB3
---------	-------

Target Details


Alternative Name:	HMGB3
Background:	HMG2A, HMG4, High mobility group protein 4
UniProt:	O15347

Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	500 µg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.



Western Blotting

Image 1. Knockout Varification: Lane 1: Wild-type A549 cell lysate, Lane 2: HMGB3 knockout A549 cell lysate, Predicted MW: 23kd Observed MW: 28kd Primary Ab: 1 µg/mL Rabbit Anti-Human HMGB3 Antibody Second Ab: 0.2 µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (ABIN6950948)