

Datasheet for ABIN7640535 **anti-Inhibin alpha antibody**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Inhibin alpha (INHA)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Inhibin alpha antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Monoclonal Antibody to Inhibin Alpha (INHa)
Immunogen:	RPA395Hu01Recombinant Inhibin Alpha (INHa)
Clone:	H2
Specificity:	The antibody is a mouse monoclonal antibody raised against INHa. It has been selected for its ability to recognize INHa in immunohistochemical staining and western blotting.
Cross-Reactivity:	Rat
Purification:	Protein A + Protein G affinity chromatography

Target Details

Target:	Inhibin alpha (INHA)
---------	----------------------

Target Details

Alternative Name:	INH _a (INHA Products)
Background:	INH-A
UniProt:	P05111
Pathways:	Peptide Hormone Metabolism , Hormone Activity , Negative Regulation of Hormone Secretion

Application Details

Application Notes:	Western blotting: 0.2-2 µg/mL, 1:500-5000 Immunohistochemistry: 5-20 µg/mL, 1:50-200 Immunocytochemistry: 5-20 µg/mL, 1:50-200 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin, Sodium azide
Precaution of Use:	This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.