

Datasheet for ABIN7640816 **anti-CD11c antibody**

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	CD11c (ITGAX)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD11c antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Monoclonal Antibody to Integrin Alpha X (CD11c)
Immunogen:	EPB159Hu61Eukaryotic Integrin Alpha X (CD11c)
Clone:	D3
Specificity:	The antibody is a mouse monoclonal antibody raised against CD11c. It has been selected for its ability to recognize CD11c in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography

Target Details

Target:	CD11c (ITGAX)
Alternative Name:	CD11c (ITGAX Products)

Target Details

Background:	ITGaX, CD11-C, ITG-AX, ITGA-X, Leu M5, Complement Component 3 Receptor 4 Subunit, CD11 antigen-like family member C, Leukocyte adhesion glycoprotein p150,95 alpha chain
UniProt:	P20702
Pathways:	Complement System , Activated T Cell Proliferation , Integrin Complex

Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.