

Datasheet for ABIN7641312

anti-IL21 Receptor antibody



Go to Product page

_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Quantity:	100 μL
Target:	IL21 Receptor (IL21R)
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL21 Receptor antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Interleukin 21 Receptor (IL21R)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against IL21R. It has been selected for its ability to recognize IL21R in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	IL21 Receptor (IL21R)
Alternative Name:	IL21R (IL21R Products)

Target Details

Background:	CD360, NILR, Novel interleukin receptor
UniProt:	Q9HBE5
Pathways:	JAK-STAT Signaling

Application Details

Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100
	Immunocytochemistry: 5-20 µg/mL,1:25-100 Optimal working dilutions must be determined by
	end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	500 μg/mL
Buffer:	PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin300.
Preservative:	Dithiothreitol (DTT), ProClin
Precaution of Use:	This product contains ProClin and Dithiothreitol (DTT): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.