

Datasheet for ABIN7641582

anti-IDH1 antibody



()	V		rV	ĺ	9	V	V
'	\mathcal{I}	٧V	<u> </u>	v	1	$\overline{}$	٧	٧

Quantity:	100 μL
Target:	IDH1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This IDH1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Target:

IDH1

Purpose:	Monoclonal Antibody to Isocitrate Dehydrogenase 1, Soluble (IDH1)
Immunogen:	RPH839Hu01Recombinant Isocitrate Dehydrogenase 1, Soluble (IDH1)
Clone:	C3
Specificity:	The antibody is a mouse monoclonal antibody raised against IDH1. It has been selected for its ability to recognize IDH1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig
Purification:	Protein A + Protein G affinity chromatography
Target Details	

Target Details

Alternative Name:	IDH1 (IDH1 Products)
Background:	PICD, IDP, Oxalosuccinate decarboxylase, NADP(+)-specific ICDH, Cytosolic NADP-isocitrate dehydrogenase
UniProt:	075874
Pathways:	Warburg Effect

Application Details

Application Notes:	Western blotting: 0.5-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-		
	20 μg/mL,Optimal working dilutions must be determined by end user.		
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated		
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious		
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration		
	date under appropriate storage condition.		
Restrictions:	For Research Use only		

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.