

Datasheet for ABIN7641710 **anti-Cytokeratin 18 antibody**

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Cytokeratin 18 (KRT18)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Cytokeratin 18 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Monoclonal Antibody to Cytokeratin 18 (CK18)
Immunogen:	RPB231Hu01Recombinant Cytokeratin 18 (CK18)
Clone:	E6
Specificity:	The antibody is a mouse monoclonal antibody raised against CK18. It has been selected for its ability to recognize CK18 in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography

Target Details

Target:	Cytokeratin 18 (KRT18)
Alternative Name:	CK18 (KRT18 Products)

Target Details

Background: KRT18, CYK18, K18, Keratin 18, Cell proliferation-inducing gene 46 protein, Keratin, type I cytoskeletal 18

UniProt: [P05783](#)

Pathways: [Apoptosis, Caspase Cascade in Apoptosis](#)

Application Details

Application Notes: Western blotting: 0.01-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Flow cytometry: 10 µg/mL, Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.