

Datasheet for ABIN7641777 **anti-KIR2DL1 antibody**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	KIR2DL1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This KIR2DL1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Monoclonal Antibody to Killer Cell Immunoglobulin Like Receptor 2DL1 (KIR2DL1)
Immunogen:	RPB509Hu01Recombinant Killer Cell Immunoglobulin Like Receptor 2DL1 (KIR2DL1)
Clone:	D11
Specificity:	The antibody is a mouse monoclonal antibody raised against KIR2DL1. It has been selected for its ability to recognize KIR2DL1 in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography

Target Details

Target:	KIR2DL1
Alternative Name:	KIR2DL1 (KIR2DL1 Products)

Target Details

Background: CD158A, CI-42, NKAT1, Killer Cell Immunoglobulin Like Receptor Two Domains, Long Cytoplasmic Tail,1, Natural killer-associated transcript 1, MHC class I NK cell receptor

UniProt: [P43626](#)

Application Details

Application Notes: Western blotting: 0.5-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.