

Datasheet for ABIN7641802 **anti-KIF5A antibody**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	KIF5A
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This KIF5A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Polyclonal Antibody to Kinesin Family, Member 5A (KIF5A)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against KIF5A. It has been selected for its ability to recognize KIF5A in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	KIF5A
Alternative Name:	KIF5A (KIF5A Products)

Target Details

Background: NKHC, MY050, SPG10, Spastic Paraplegia 10(Autosomal Dominant), Kinesin heavy chain neuron-specific 1, Neuronal kinesin heavy chain

UniProt: [Q6QLM7](#)

Application Details

Application Notes: Western blotting: 0.2-2 µg/mL, 1:250-2500 Immunohistochemistry: 5-20 µg/mL, 1:25-100 Immunocytochemistry: 5-20 µg/mL, 1:25-100 Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 500 µg/mL

Buffer: PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin300.

Preservative: Dithiothreitol (DTT), ProClin

Precaution of Use: This product contains ProClin and Dithiothreitol (DTT): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.