

## Datasheet for ABIN7641950

## anti-LAMB2 antibody



_					
	W	0	rv	10	W

Quantity:	100 μL	
Target:	LAMB2	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This LAMB2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

## **Product Details**

Alternative Name:

Background:

Purpose:	Polyclonal Antibody to Laminin Beta 2 (LAMb2)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against LAMb2. It has been selected for its ability to recognize LAMb2 in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	
Target Details		
Target:	LAMB2	

LAM-B2, LAMS, Laminin S, Laminin B1s chain, S-laminin subunit beta

LAMb2 (LAMB2 Products)

## **Target Details**

UniProt:	P55268	
Pathways:	Skeletal Muscle Fiber Development	
Application Dataila		
Application Details		
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100	
	Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by	
	end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	500 μg/mL	
Buffer:	PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin300.	
Preservative:	Dithiothreitol (DTT), ProClin	
Precaution of Use:	This product contains ProClin and Dithiothreitol (DTT): POISONOUS AND HAZARDOUS	
	SUBSTANCES which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	