

Datasheet for ABIN7642039

anti-LMAN1 antibody



Overview

Quantity:	100 μL
Target:	LMAN1
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LMAN1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC), Flow Cytometry (FACS)

Product Details

Target:

Alternative Name:

Purpose:	Polyclonal Antibody to Lectin, Mannose Binding 1 (LMAN1)
Immunogen:	RPC567Hu01Recombinant Lectin, Mannose Binding 1 (LMAN1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against LMAN1. It has been selected for its ability to recognize LMAN1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

LMAN1

LMAN1 (LMAN1 Products)

Target Details

Background:	F5F8D, FMFD1, MCFD1, MR60, Gp58, Protein ERGIC-53, Coagulation Factor V-Factor VIII
	Combined Deficiency, Endoplasmic Reticulum-Golgi Intermediate Compartment Protein 53
UniProt:	P49257
Application Details	
Application Notes:	Western blotting: 0.01-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunofluorescence:5-20 μg/mL,Flow cytometry:20 μg/mL,Optimal working dilutions must be determined by end user.,
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.