

## Datasheet for ABIN7642049

## anti-LEFTY2 antibody



Go to Product page

_					
	W	0	rv	10	W

Quantity:	100 μL	
Target:	LEFTY2	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This LEFTY2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

## **Product Details**

Purpose:	Polyclonal Antibody to Left/Right Determination Factor 2 (LEFTY2)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against LEFTY2. It has been selected for its ability to recognize LEFTY2 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

Target:	LEFTY2
Alternative Name:	LEFTY2 (LEFTY2 Products)

## **Target Details**

Background:	TGFB4, EBAF, LEFTA, LEFTYA, Endometrial Bleeding Associated Factor, Left-right	
	determination factor A, Transforming growth factor beta-4	
UniProt:	Q5UCE3	
Application Details		
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	500 μg/mL	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	