

#### Datasheet for ABIN7642917

# anti-MOSC1 antibody



()	ve	r\/i		۱۸/
$\cup$	V C	1 / 1	$\overline{}$	٧V

Quantity:	100 μL
Target:	MOSC1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This MOSC1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

### **Product Details**

Purpose:	Monoclonal Antibody to MOCO Sulphurase C-Terminal Domain Containing Protein 1 (MOSC1)
Specificity:	The antibody is a mouse monoclonal antibody raised against MOSC1. It has been selected for its ability to recognize MOSC1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

## **Target Details**

Target:	MOSC1
Alternative Name:	MOSC1 (MOSC1 Products)
Background:	Molybdenum Cofactor Sulphurase C-Terminal Domain Containing Protein, Mitochondrial amidoxime-reducing component 1
UniProt:	Q5VT66

## **Target Details**

Pathways:	SARS-CoV-2 Protein Interactome	
Application Details		
Application Notes:	Western blotting: 0.2-2 μg/mL,1:500-5000 Immunohistochemistry: 5-20 μg/mL,1:50-200 Immunocytochemistry: 5-20 μg/mL,1:50-200 Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	