

Datasheet for ABIN7643232

anti-MPG antibody



Overview

Quantity:	100 μL
Target:	MPG
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MPG antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Target:

Alternative Name:

MPG

Purpose:	Polyclonal Antibody to N-Methylpurine DNA Glycosylase (MPG)
Immunogen:	RPH625Ra01Recombinant NMethylpurine DNA Glycosylase (MPG)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against MPG. It has been selected for its ability to recognize MPG in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

N-Methylpurine DNA Glycosylase (MPG Products)

Target Details

Target Details	
Background:	AAG, APNG, ADPG, CRA36.1, MDG, Mid1, PIG11, PIG16, Anpg, DNA-3-Methyladenine
	Glycosylase, Alkyladenine DNA Glycosylase, 3-alkyladenine DNA glycosylase
UniProt:	P23571
Pathways:	DNA Damage Repair
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL,lmmunohistochemistry: 5-20 μg/mL,lmmunocytochemistry: 5-
	20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	500 μg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without

detectable loss of activity. Avoid repeated freeze-thaw cycles.