

Datasheet for ABIN7643288

anti-NCK1 antibody



Overview

Quantity:	100 μL
Target:	NCK1
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NCK1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Cytoplasmic Protein NCK1 (NCK1)
Immunogen:	RPC712Hu01Recombinant Cytoplasmic Protein NCK1 (NCK1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against NCK1. It has been selected for its ability to recognize NCK1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	NCK1

Target Details

rarget Details	
Alternative Name:	NCK1 (NCK1 Products)
Background:	NCKalpha, NCKa, NCK adaptor protein 1, Non Catalytic Region Of Tyrosine Kinase Adaptor
	Protein 1, SH2/SH3 adaptor protein NCK-alpha
UniProt:	P16333
Pathways:	TCR Signaling, Regulation of Actin Filament Polymerization, Signaling Events mediated by
	VEGFR1 and VEGFR2, VEGFR1 Specific Signals
Application Details	
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100
	Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by
	end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
	date diluei appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	500 μg/mL
Buffer:	PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin300.
Preservative:	Dithiothreitol (DTT), ProClin
Precaution of Use:	This product contains ProClin and Dithiothreitol (DTT): POISONOUS AND HAZARDOUS
	SUBSTANCES which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.