

Datasheet for ABIN7643383

anti-Neuregulin 4 antibody



Overview

Quantity:	100 μL
Target:	Neuregulin 4 (NRG4)
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Neuregulin 4 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Target:

Purpose:	Polyclonal Antibody to Neuregulin 4 (NRG4)
Immunogen:	RPC174Mu01Recombinant Neuregulin 4 (NRG4)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against NRG4. It has been selected for its ability to recognize NRG4 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

Neuregulin 4 (NRG4)

Target Details

Alternative Name:	NRG4 (NRG4 Products)
Background:	HRG4, Pro-NRG4, Pro-neuregulin-4, membrane-bound isoform
UniProt:	Q9WTX4
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway

Application Details

Application Notes:	Western blotting: 0.5-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-
	20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.78 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.