

Datasheet for ABIN7643723

anti-Glucocorticoid Receptor antibody





Go to Product page

_				
()	ve.	rv/	101	Λ

Quantity:100 μLTarget:Glucocorticoid Receptor (NR3C1)Reactivity:HumanHost:RabbitClonality:Polyclonal
Reactivity: Human Host: Rabbit
Host: Rabbit
Clonality: Polyclonal
Conjugate: This Glucocorticoid Receptor antibody is un-conjugated
Application: Western Blotting (WB), Immunohistochemistry (IHC)
Product Details

Product Details


Purpose:	Polyclonal Antibody to Glucocorticoid Receptor (GR)	
Immunogen:	RPB608Hu02Recombinant Glucocorticoid Receptor (GR)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against GR. It has been selected for its ability to recognize GR in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	

Target Details

Target:	Glucocorticoid Receptor (NR3C1)
Alternative Name:	Glucocorticoid Receptor
Background:	GCCR, GCR, NR3C1, GRL, NR3-C1, Nuclear Receptor Subfamily 3, Group C, Member 1

Target Details

rarger Details		
UniProt:	P04150	
Pathways:	Nuclear Receptor Transcription Pathway, Intracellular Steroid Hormone Receptor Signaling	
	Pathway, Steroid Hormone Mediated Signaling Pathway, Regulation of Intracellular Steroid	
	Hormone Receptor Signaling, Regulation of Hormone Metabolic Process, Regulation of	
	Hormone Biosynthetic Process, Regulation of Muscle Cell Differentiation, Regulation of	
	Carbohydrate Metabolic Process	
Application Details		
Application Notes:	Western blotting: 0.5-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-	
	20 μg/mL,Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	0.5 mg/mL	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be	
	handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	



Western Blotting

Image 1. Knockout Varification: Lane 1: Wild-type HepG2 cell lysate, Lane 2: NR3C1 knockout HepG2 cell lysate, Predicted MW: 78~85,60~64kd Observed MW: 97kd Primary Ab: 2 μg/mL Rabbit Anti-Human NR3C1 Antibody Second Ab: 0.2 μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (ABIN6950948)