

### Datasheet for ABIN7643729

# anti-Glucocorticoid Receptor antibody



#### Go to Product page

_				
( )	ve.	rv/	101	Λ

Quantity:	100 μL	
Target:	Glucocorticoid Receptor (NR3C1)	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This Glucocorticoid Receptor antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

### **Product Details**

**Target Details** 

Purpose:	Monoclonal Antibody to Glucocorticoid Receptor (GR)	
Immunogen:	RPB608Hu03Recombinant Glucocorticoid Receptor (GR)	
Clone:	C8	
Specificity:	The antibody is a mouse monoclonal antibody raised against GR. It has been selected for its ability to recognize GR in immunohistochemical staining and western blotting.	
Purification:	Protein A + Protein G affinity chromatography	

Target: Glucocorticoid Receptor (NR3C1)

Alternative Name: Glucocorticoid Receptor (NR3C1 Products)

## Target Details

Background:	GCCR, GCR, NR3C1, GRL, NR3-C1, Nuclear Receptor Subfamily 3, Group C, Member 1	
UniProt:	P04150	
Pathways:	Nuclear Receptor Transcription Pathway, Intracellular Steroid Hormone Receptor Signaling Pathway, Steroid Hormone Mediated Signaling Pathway, Regulation of Intracellular Steroid Hormone Receptor Signaling, Regulation of Hormone Metabolic Process, Regulation of Hormone Biosynthetic Process, Regulation of Muscle Cell Differentiation, Regulation of Carbohydrate Metabolic Process	

	Hormone Receptor Signaling, Regulation of Hormone Metabolic Process, Regulation of Hormone Biosynthetic Process, Regulation of Muscle Cell Differentiation, Regulation of Carbohydrate Metabolic Process	
Application Details		
Application Notes:	Western blotting: 0.2-2 μg/mL,1:500-5000 Immunohistochemistry: 5-20 μg/mL,1:50-200 Immunocytochemistry: 5-20 μg/mL,1:50-200 Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions: Handling	For Research Use only	
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.	
Preservative:	ProClin, Sodium azide	
Precaution of Use:	This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	