

Datasheet for ABIN7643764

anti-NUP153 antibody



Overview

| Quantity: | 100 μL |
|--------------|--|
| Target: | NUP153 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This NUP153 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details

| tigen-specific affinity chromatography followed by Protein A affinity chromatography |
|--|
| |
| t |
| |
| e antibody is a rabbit polyclonal antibody raised against NUP153. It has been selected for its ility to recognize NUP153 in immunohistochemical staining and western blotting. |
| |
| B147Hu01Recombinant Nucleoporin 153 (NUP153) |
| lyclonal Antibody to Nucleoporin 153 (NUP153) |
| |

| Target Details | |
|---------------------|---|
| Alternative Name: | NUP153 (NUP153 Products) |
| Background: | HNUP153, N153, Nucleoporin 153 kDa |
| UniProt: | P49790 |
| Pathways: | Protein targeting to Nucleus |
| Application Details | |
| Application Notes: | Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 0.5 mg/mL |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. |
| Preservative: | ProClin, Sodium azide |