

Datasheet for ABIN7643823
anti-OPA3 antibody



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	OPA3
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OPA3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Optic Atrophy 3 (OPA3)
Immunogen:	RPC701Ra01Recombinant Optic Atrophy 3 (OPA3)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against OPA3. It has been selected for its ability to recognize OPA3 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	OPA3
---------	------

Target Details

Alternative Name:	OPA3 (OPA3 Products)
Background:	MGA3, Recessive,With Chorea And Spastic Paraplegia, 3-Methylglutaconic Aciduria Type 3
UniProt:	B5DF67

Application Details

Application Notes:	Western blotting: 0.2-2 µg/mL, 1:250-2500 Immunohistochemistry: 5-20 µg/mL, 1:25-100 Immunocytochemistry: 5-20 µg/mL, 1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	500 µg/mL
Buffer:	PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin300.
Preservative:	Dithiothreitol (DTT), ProClin
Precaution of Use:	This product contains ProClin and Dithiothreitol (DTT): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.