

## Datasheet for ABIN7644234

## anti-Phenylalanine Hydroxylase antibody



()	11/	$\sim$	r١.	ıi.	0	۱۸/	,
U	V	H	r٧	1	C	V۷	

Quantity:	100 μL
Target:	Phenylalanine Hydroxylase
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Phenylalanine Hydroxylase antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

## **Product Details**

Target:	Phanylalanina Hydroxylasa
Target Details	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Specificity:	The antibody is a rabbit polyclonal antibody raised against PAH. It has been selected for its ability to recognize PAH in immunohistochemical staining and western blotting.
Isotype:	IgG
Purpose:	Polyclonal Antibody to Phenylalanine Hydroxylase (PAH)
Durana a	Delvoland Antibady to Diagnodalania dividuanda a (DALI)

Target:	Phenylalanine Hydroxylase
Abstract:	Phenylalanine Hydroxylase Products
Background:	PKU, PKU1, Phe-4-monooxygenase

## **Target Details**

UniProt:	P16331
Application Details	
Application Notes:	Western blotting: 0.01-2 $\mu$ g/mL,Immunohistochemistry: 5-20 $\mu$ g/mL,Immunocytochemistry: 5-20 $\mu$ g/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.