

### Datasheet for ABIN7644288

# anti-PGAM1 antibody



#### Overview

Quantity:	100 μL
Target:	PGAM1
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This PGAM1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

### **Product Details**

Purpose:	Monoclonal Antibody to Phosphoglycerate Mutase 1, Brain (PGAM1)
Clone:	D11
Specificity:	The antibody is a mouse monoclonal antibody raised against PGAM1. It has been selected for its ability to recognize PGAM1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Rat
Purification:	Protein A + Protein G affinity chromatography
Target Detaile	

#### Target Details

Target:	PGAM1
Alternative Name:	PGAM1 (PGAM1 Products)

## **Target Details**

rarget Details	
Background:	PGAMA, PGAM-B, BPG-dependent PGAM 1, Phosphoglycerate mutase isozyme B
UniProt:	P18669
Pathways:	Regulation of Carbohydrate Metabolic Process
Application Details	
Application Notes:	Western blotting: $0.2-2~\mu g/m L$ , $1:500-5000~lmmunohistochemistry$ : $5-20~\mu g/m L$ , $1:50-200~lmmunocytochemistry$ : $5-20~\mu g/m L$ , $1:50-200~optimal~working~dilutions~must~be~determined~by~end~user$ .
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS, pH 7.4, containing 0.01 % SKL, 1 mM DTT, 5 % Trehalose and Proclin300.
Preservative:	Dithiothreitol (DTT), ProClin
Precaution of Use:	This product contains ProClin and Dithiothreitol (DTT): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.