

Datasheet for ABIN7644296

anti-PGAM2 antibody



()	ve	r\/i	Δ	۱۸/
\circ	V C	1 V		v v

Quantity:	100 μL	
Target:	PGAM2	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PGAM2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Purpose:	Polyclonal Antibody to Phosphoglycerate Mutase 2, Muscle (PGAM2)	
Immunogen:	RPB645Hu01Recombinant Phosphoglycerate Mutase 2, Muscle (PGAM2)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against PGAM2. It has been selected for its ability to recognize PGAM2 in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Mouse, Rat	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	
Target Details		
Target:	PGAM2	

Target Details

end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelera thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expirat date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANI which should be handled by trained staff only. Storage: 4 °C,-20 °C	. d g d t 2 d t d d		
Phosphoglycerate mutase isozyme M UniProt: P15259 Application Details Application Notes: Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelera thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expirat date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTAN which should be handled by trained staff only. Storage: 4 °C, 20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without the staff only.	Alternative Name:	PGAM2 (PGAM2 Products)	
UniProt: P15259 Application Details Application Notes: Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerate thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expirate date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTAN which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without the process of the	Background:	PGAM-M, BPG-dependent PGAM 2, Muscle-specific phosphoglycerate mutase,	
Application Details Application Notes: Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerathermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expirated ate under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANI which should be handled by trained staff only. Storage: 4 °C, -20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without the staff on the		Phosphoglycerate mutase isozyme M	
Application Notes: Western blotting: 0.2-2 µg/mL,1:250-2500 Immunohistochemistry: 5-20 µg/mL,1:25-100 Immunocytochemistry: 5-20 µg/mL,1:25-100 Optimal working dilutions must be determined end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerate thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expirate date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANI which should be handled by trained staff only. Storage: 4 *C,-20 *C Storage Comment: Store at 4*C for frequent use. Stored at -20*C in a manual defrost freezer for two year without the process of the proc	UniProt:	P15259	
Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelera thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expirat date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTAN which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without the staff on the contains	Application Details		
end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelera thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expirat date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANI which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without the storage of the proclin and sodium and proclin a manual defrost freezer for two year without the storage of the proclin and sodium and so the proclin and sodium azide.	Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100	
thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expirat date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANG which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without the storage of the process of the proc		Immunocytochemistry: 5-20 μ g/mL,1:25-100 Optimal working dilutions must be determined by end user.	
degradation and precipitation were observed. The loss rate is less than 5% within the expirate date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANG which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without the staff of the s	Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
date under appropriate storage condition. Restrictions: For Research Use only Handling Format: Liquid Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANG which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without the storage of the storag		thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
Restrictions: Handling Format: Liquid Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANG which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without the storage of the stor		degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
Format: Liquid Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANG which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without		date under appropriate storage condition.	
Format: Liquid Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANG which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	Restrictions:	For Research Use only	
Concentration: 0.5 mg/mL Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANG which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	Handling		
Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANG which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	Format:	Liquid	
Preservative: ProClin, Sodium azide Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANG which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	Concentration:	0.5 mg/mL	
Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANG which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
which should be handled by trained staff only. Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	Preservative:	ProClin, Sodium azide	
Storage: 4 °C,-20 °C Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	Precaution of Use:	This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANCES	
Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without		which should be handled by trained staff only.	
·	Storage:	4 °C,-20 °C	
	Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.	