

Datasheet for ABIN7644504

anti-PDGFA antibody



Overview

Quantity:	100 μL
Target:	PDGFA
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This PDGFA antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Monoclonal Antibody to Platelet Derived Growth Factor Subunit A (PDGFA)
Immunogen:	RPA528Hu01Recombinant Platelet Derived Growth Factor Subunit A (PDGFA)
Clone:	C4
Specificity:	The antibody is a mouse monoclonal antibody raised against PDGFA. It has been selected for its ability to recognize PDGFA in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography

Target Details

Target:	PDGFA
Alternative Name:	PDGFA (PDGFA Products)

Target Details	
Background:	PDGF-A, PDGF1, Platelet Derived Growth Factor Alpha Polypeptide
UniProt:	P04085
Pathways:	RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Smooth Muscle Cell Migration, Platelet-derived growth Factor Receptor Signaling
Application Details	
Application Notes:	Western blotting: 0.01-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.