

Datasheet for ABIN7644607

anti-PKD2 antibody



| (| ۱۱/ | e | r\/ | Ì١ | | ۱۸ | |
|---|-----|--------|-----|----|---------------|----|---|
| | , v | \cup | V | 1 | $\overline{}$ | V | V |

| Quantity: | 100 μL |
|--------------|--|
| Target: | PKD2 |
| Reactivity: | Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PKD2 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details

| Product Details | |
|-------------------|--|
| Purpose: | Polyclonal Antibody to Protein Kinase D2 (PKD2) |
| Immunogen: | RPA510Mu01Recombinant Protein Kinase D2 (PKD2) |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against PKD2. It has been selected for its ability to recognize PKD2 in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Human, Rat |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |
| Target Details | |
| Target: | PKD2 |

Target Details

| Alternative Name: | PKD2 (PKD2 Products) |
|-------------------|---|
| Background: | PRKD2, PRK-D2, PRKD-2, PK-D2, HSPC187, nPKC-D2, Serine/threonine-protein kinase D2 |
| UniProt: | Q8BZ03 |
| Pathways: | cAMP Metabolic Process, Maintenance of Protein Location, Negative Regulation of Transporter Activity |

Application Details

| Application Notes: | Western blotting: 0.5-5 μg/mL,Immunohistochemistry: 5-50 μg/mL,Immunocytochemistry: 5-50 μg/mL,Optimal working dilutions must be determined by end user. |
|--------------------|--|
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |
| | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious |
| | degradation and precipitation were observed. The loss rate is less than 5% within the expiration |
| | date under appropriate storage condition. |
| Restrictions: | For Research Use only |

Handling

| Format: | Liquid |
|--------------------|---|
| Concentration: | 500 μg/mL |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. |