

Datasheet for ABIN7644661

anti-Melanoma gp100 antibody



Overview

Quantity:	100 μL
Target:	Melanoma gp100 (PMEL)
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Melanoma gp100 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Melanoma Associated ME20 (ME20M)
Immunogen:	RPB330Hu01Recombinant Melanoma Associated ME20 (ME20M)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against ME20M. It has been selected for its ability to recognize ME20M in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

Target:	Melanoma gp100 (PMEL)
Alternative Name:	ME20M (PMEL Products)

Target Details

Background:	PMEL, GP100, SI, SIL, SILV, Pmel17, Silver Homolog, Premelanosome Protein, Silver Locus
	Protein Homolog, 95 kDa melanocyte-specific secreted glycoprotein
UniProt:	P40967
Application Details	
Application Notes:	Western blotting: 0.01-2 μg/mL,lmmunohistochemistry: 5-20 μg/mL,lmmunocytochemistry: 5-
	20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.