

Datasheet for ABIN7644710 **anti-CXCL7 antibody**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	CXCL7 (PPBP)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CXCL7 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Monoclonal Antibody to Chemokine (C-X-C motif) ligand 7 (CXCL7)
Immunogen:	RPA370Hu01Recombinant Chemokine (CXC motif) ligand 7 (CXCL7)
Clone:	C5
Specificity:	The antibody is a mouse monoclonal antibody raised against CXCL7. It has been selected for its ability to recognize CXCL7 in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography

Target Details

Target:	CXCL7 (PPBP)
Alternative Name:	CXCL7 (PPBP Products)

Target Details

Background: CXCL7, PPBP, PBP, B-TG1, CTAP-III, CTAP3, CTAPIII, LA-PF4, LDGF, MDGF, NAP2, SCYB7, TC1, TC2, TGB1, THBGB1, Pro-Platelet Basic Protein, Beta-Thromboglobulin

UniProt: [P02775](#)

Application Details

Application Notes: Western blotting: 0.2-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.

Preservative: ProClin, Sodium azide

Precaution of Use: This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.