

Datasheet for ABIN7644856
anti-PIP antibody



[Go to Product page](#)

Overview

| | |
|--------------|--------------------------------------------------------------------------------------------------------|
| Quantity: | 100 µL |
| Target: | PIP |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PIP antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP) |

Product Details

| | |
|---------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Purpose: | Polyclonal Antibody to Prolactin Induced Protein (PIP) |
| Immunogen: | RPM035Ra01 Recombinant Prolactin Induced Protein (PIP) |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against PIP. It has been selected for its ability to recognize PIP in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |

Target Details

| | |
|-------------------|------------------------------------------------------------|
| Target: | PIP |
| Alternative Name: | Prolactin Induced Protein (PIP Products) |

Target Details

| | |
|-------------|--------------------------------------------------------------------------------------------------------------------------------------|
| Background: | GCDFP15, GPIP4, BRST2, SABP, gp17, Gross cystic disease fluid protein 15, Prolactin-induced protein, Secretory actin-binding protein |
| UniProt: | O70417 |

Application Details

| | |
|--------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Application Notes: | Western blotting: 0.01-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunofluorescence: 5-20 µg/mL, Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |

Handling

| | |
|--------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Format: | Liquid |
| Concentration: | 0.5 mg/mL |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C, -20 °C |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. |