

Datasheet for ABIN7644902 **anti-POMC antibody**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	POMC
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This POMC antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Polyclonal Antibody to Proopiomelanocortin (POMC)
Immunogen:	RPB311Mu01Recombinant Proopiomelanocortin (POMC)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against POMC. It has been selected for its ability to recognize POMC in immunohistochemical staining and western blotting.
Cross-Reactivity:	Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details

Target:	POMC
---------	------

Target Details

Alternative Name:	POMC (POMC Products)
Background:	POC, ProOMC, Pro-OMC, Pro Opiomelanocortin, Corticotropin-lipotropin
UniProt:	P01193
Pathways:	Metabolism of Steroid Hormones and Vitamin D , Peptide Hormone Metabolism , Hormone Activity , Feeding Behaviour

Application Details

Application Notes:	Western blotting: 0.5-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	500 µg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.