

Datasheet for ABIN7644908

anti-PCCB antibody



\sim				
	1//	Д	rv	۱۸/

Quantity:	100 μL	
Target:	PCCB	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PCCB antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Purpose:	Polyclonal Antibody to Propionyl Coenzyme A Carboxylase Beta (PCCb)		
Isotype:	IgG		
Specificity:	The antibody is a rabbit polyclonal antibody raised against PCCb. It has been selected for its ability to recognize PCCb in immunohistochemical staining and western blotting.		
Cross-Reactivity:	Human, Rat		
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography		
Target Details			

Target:	PCCB
Alternative Name:	PCCb (PCCB Products)

Target Details

Background:	Propanoyl-CoA:carbon dioxide ligase subunit beta, Propionyl-CoA carboxylase beta chain,	
	mitochondrial	
UniProt:	Q7TMF4	
Pathways:	Monocarboxylic Acid Catabolic Process	
Application Details		
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100	
	Immunocytochemistry: 5-20 μ g/mL,1:25-100 Optimal working dilutions must be determined by	
	end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	500 μg/mL	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
	should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
	detectable loss of activity. Avoid repeated freeze-thaw cycles.	