

Datasheet for ABIN7644926

anti-Prostacyclin Receptor antibody



Overview

Quantity:	100 μL
Target:	Prostacyclin Receptor (PTGIR)
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Prostacyclin Receptor antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Alternative Name:

Background:

Purpose:	Polyclonal Antibody to Prostacyclin Receptor (PTGIR)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PTGIR. It has been selected for its ability to recognize PTGIR in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	Prostacyclin Receptor (PTGIR)

PTGI-R, IP1, PRIPR, Prostaglandin I2 Receptor, Prostanoid IP receptor

PTGIR (PTGIR Products)

Target Details

UniProt:	P43119
Pathways:	cAMP Metabolic Process, Platelet-derived growth Factor Receptor Signaling, Thromboxane A2 Receptor Signaling
Application Details	
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100 Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions: Handling	For Research Use only
Format:	Liquid
Concentration:	500 μg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.