

Datasheet for ABIN7645054

anti-PKC gamma antibody



Overview

| Quantity: | 100 μL |
|--------------|--|
| Target: | PKC gamma (PRKCG) |
| Reactivity: | Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This PKC gamma antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details

Target:

| - Toddet Details | |
|-------------------|--|
| Purpose: | Polyclonal Antibody to Protein Kinase C Gamma (PKCg) |
| Immunogen: | RPA419Ra01Recombinant Protein Kinase C Gamma (PKCg) |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against PKCg. It has been selected for its ability to recognize PKCg in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Human, Mouse |
| Purification: | Antigen-specific affinity chromatography followed by Protein A affinity chromatography |
| Target Details | |

PKC gamma (PRKCG)

Target Details

| Alternative Name: | PKCg (PRKCG Products) |
|-------------------|---|
| Background: | PKC-G, PRKCG, PKCC, SCA14, PKC-Gamma |
| UniProt: | P63319 |
| Pathways: | WNT Signaling, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Thyroid Hormone Synthesis, Myometrial Relaxation and Contraction, G-protein mediated Events, Positive Regulation of Response to DNA Damage Stimulus, Interaction of EGFR with phospholipase C-gamma, Thromboxane A2 Receptor Signaling, VEGF Signaling |

Application Details

| Application Notes: | Western blotting: 0.5-2 μg/mL,lmmunohistochemistry: 5-20 μg/mL,lmmunocytochemistry: 5-20 μg/mL,Optimal working dilutions must be determined by end user. |
|--------------------|---|
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |

| Format: | Liquid |
|--------------------|---|
| Concentration: | 500 μg/mL |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. |