

Datasheet for ABIN7645071

anti-PRKAG2 antibody



Go to Product page

_						
	V	\triangle	r۱	/1	\triangle	Λ/
	' V '		ΙV			v v

Quantity:	100 μL	
Target:	PRKAG2	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This PRKAG2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Purpose:	Polyclonal Antibody to Protein Kinase, AMP Activated Gamma 2 (PRKAg2)
Immunogen:	RPD356Mu01Recombinant Protein Kinase, AMP Activated Gamma 2 (PRKAg2)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PRKAg2. It has been selected for its ability to recognize PRKAg2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

Target:	PRKAG2
Alternative Name:	PRKAg2 (PRKAG2 Products)

Target Details	
Background:	AAKG, AAKG2, CMH6, H91620p, WPWS, 5'-AMP-activated protein kinase subunit gamma-2
UniProt:	Q91WG5
Pathways:	AMPK Signaling, Cellular Glucan Metabolic Process, Ribonucleoside Biosynthetic Process, Regulation of Carbohydrate Metabolic Process, Warburg Effect
Application Details	
Application Notes:	Western blotting: 0.01-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.