

Datasheet for ABIN7645072

anti-PRKAG3 antibody



Overview

Quantity:	100 μL
Target:	PRKAG3
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PRKAG3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Target:

Alternative Name:

Purpose:	Polyclonal Antibody to Protein Kinase, AMP Activated Gamma 3 (PRKAg3)
Immunogen:	RPG288Mu01Recombinant Protein Kinase, AMP Activated Gamma 3 (PRKAg3)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PRKAg3. It has been selected for its ability to recognize PRKAg3 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

PRKAG3

PRKAg3 (PRKAG3 Products)

Target Details

Background:	AMPKG3, 5'-AMP-activated protein kinase subunit gamma-3
UniProt:	Q8BGM7
Pathways:	AMPK Signaling, Cellular Glucan Metabolic Process, Warburg Effect
Application Details	
Application Notes:	Western blotting: 0.2-2 μg/mL,1:250-2500 Immunohistochemistry: 5-20 μg/mL,1:25-100
	Immunocytochemistry: 5-20 μg/mL,1:25-100 Optimal working dilutions must be determined by
	end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.45 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.