

### Datasheet for ABIN7645132

# anti-PTPRG antibody



#### Overview

Quantity:	100 μL
Target:	PTPRG
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PTPRG antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

#### **Product Details**

FIOUUCI Details	
Purpose:	Polyclonal Antibody to Protein Tyrosine Phosphatase Receptor Type G (PTPRG)
Immunogen:	RPD602Hu01Recombinant Protein Tyrosine Phosphatase Receptor Type G (PTPRG)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PTPRG. It has been selected for its ability to recognize PTPRG in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	PTPRG

## **Target Details**

raiget Details	
Alternative Name:	PTPRG (PTPRG Products)
Background:	PTPR-G, HPTPG, PTPG, R-PTP-GAMMA, RPTPG, Receptor-type tyrosine-protein phosphatase
	gamma
UniProt:	P23470
Application Details	
Application Notes:	Western blotting: 0.01-2 μg/mL,lmmunohistochemistry: 5-20 μg/mL,lmmunocytochemistry: 5-
	20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.