

Datasheet for ABIN7645237

anti-PDP2 antibody



Overview

Quantity:	100 μL
Target:	PDP2
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PDP2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Polyclonal Antibody to Pyruvate Dehydrogenase Phosphatase 2 (PDP2)
Immunogen:	RPD743Hu01Recombinant Pyruvate Dehydrogenase Phosphatase 2 (PDP2)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PDP2. It has been selected for its ability to recognize PDP2 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	
Target:	PDP2

Target Details

ranget betails	
Alternative Name:	PDP2 (PDP2 Products)
Background:	PPM2C2, PDPC 2, Protein Phosphatase 2C, Magnesium-Dependent, Catalytic Subunit 2,
	[Pyruvate dehydrogenase [acetyl-transferring]]-phosphatase 2, mitochondrial
UniProt:	Q9P2J9
Application Details	
Application Notes:	Western blotting: 0.01-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-
	20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.