

Datasheet for ABIN7645337 **anti-REG1A antibody**

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	REG1A
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This REG1A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Monoclonal Antibody to Regenerating Islet Derived Protein 1 Alpha (REG1a)
Immunogen:	RPB760Hu01Recombinant Regenerating Islet Derived Protein 1 Alpha (REG1a)
Clone:	C7
Specificity:	The antibody is a mouse monoclonal antibody raised against REG1a. It has been selected for its ability to recognize REG1a in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography

Target Details

Target:	REG1A
Alternative Name:	REG1a (REG1A Products)

Target Details

Background: REG, PSP, ICRF, PSPS1, PTP, Islet cells regeneration factor, Islet of Langerhans regenerating protein, Lithostathine 1 Alpha, Pancreatic Stone Protein, Pancreatic Thread Protein

UniProt: [P05451](#)

Application Details

Application Notes: Western blotting: 0.01-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.