

Datasheet for ABIN7645443

anti-RETNLB antibody



Overview

Quantity:	100 μL
Target:	RETNLB
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RETNLB antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Target:

Alternative Name:

Purpose:	Polyclonal Antibody to Resistin Like Beta (RETNLb)
Immunogen:	RPM131Hu01Recombinant Resistin Like Beta (RETNLb)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against RETNLb. It has been selected for its ability to recognize RETNLb in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography
Target Details	

RETNLB

RETNLb (RETNLB Products)

Target Details

Target Details	
Background:	XCP2, FIZZ2, HXCP2, CCRG, RETNL2, RELMb, Foundin In Flammatory Zone 2, Colon and small
	intestine-specific cysteine-rich protein, Colon carcinoma-related gene protein
UniProt:	Q9BQ08
Pathways:	Hormone Activity
Application Details	
Application Notes:	Western blotting: 0.01-2 μg/mL,Immunohistochemistry: 5-20 μg/mL,Immunocytochemistry: 5-
	20 μg/mL,Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	0.29 mg/mL
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.