

Datasheet for ABIN7646163

anti-STAT2 antibody



_					
	1//	r	Vİ	\triangle	۸/
	V		VI		/ V

Quantity:	100 μL	
Target:	STAT2	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This STAT2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)	

Product Details

Purpose:	Polyclonal Antibody to Signal Transducer And Activator Of Transcription 2 (STAT2)	
Immunogen:	RPB796Mu01Recombinant Signal Transducer And Activator Of Transcription 2 (STAT2)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against STAT2. It has been selected for its ability to recognize STAT2 in immunohistochemical staining and western blotting.	
Cross-Reactivity:	Human	
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography	
Target Details		
Target:	STAT2	

Target Details

Storage Comment:

l arget Details		
Alternative Name:	STAT2 (STAT2 Products)	
Background:	ISGF-3, P113, STAT113	
UniProt:	Q9WVL2	
Pathways:	JAK-STAT Signaling, Hepatitis C, CXCR4-mediated Signaling Events	
Application Details		
Application Notes:	Western blotting: 0.2-2 μ g/mL,1:250-2500 Immunohistochemistry: 5-20 μ g/mL,1:25-100 Immunocytochemistry: 5-20 μ g/mL,1:25-100 Optimal working dilutions must be determined by end user.	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	0.5 mg/mL	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Preservative:	ProClin, Sodium azide	
Precaution of Use:	This product contains ProClin and Sodium azide: POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	

detectable loss of activity. Avoid repeated freeze-thaw cycles.

Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without